



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

progression; or that the sum of their squares shall be a square, or that they shall be the sides of a right-angled triangle. In fact the number of requirements is unlimited.

In the first case, let  $a = 2rs - r^2 + s^2$ ,  $b = r^2 + s^2$ ,  $c = 2rs + r^2 - s^2$ , which are general expressions for three square numbers in arithmetical progression;  $r$ ,  $s$ , being any different numbers that will make  $a$ ,  $b$ ,  $c$ , all positive. Let  $r = 2$ ,  $s = 1$ ; then  $a = 1$ ,  $b = 5$ ,  $c = 7$ , and

$$x = \frac{151321}{7863240} \dots \frac{151321}{7863240}, \frac{756605}{7863240}, \text{ and } \frac{1059247}{7863240}$$

are the roots for the positive sign in  $a^2x^2 \pm ax = \square$ , &c. In order to have a positive value of  $x$  to satisfy the negative sign, take  $r = 4$ ,  $s = 3$ ;

then  $a=17$ ,  $b=25$ ,  $c=31$ , and  $x = \frac{(864571)^2}{11011044931800}$ . Therefore

$$\frac{12707211238697}{11011044931800}, \frac{18687075351025}{11011044931800}, \frac{23171973435271}{11011044931800}$$

are the roots for the negative sign in  $a^2x^2 \pm ax = \square$ , &c.

## SOLUTIONS OF PROBLEMS IN NUMBER 2.

101.—“From any point  $O$ , within the circumference of a circle, two lines are drawn making a constant angle with each other. These lines revolve about  $O$  in the plane of the circle, and from the points where they cut the circumference tangents are drawn. Find the locus of the intersection of these tangents.

SOLUTION BY PROF. W. W. BEMAN, ANN ARBOR, MICHIGAN.

LET a diameter through  $O$  represent the axis of  $X$ . Put  $r$  = the radius of the given circle,  $a$  = the distance of the point  $O$  from its centre,  $\alpha$  = the constant angle included by the two given lines;  $x, y$ , coordinates of the intersection of the tangents, the origin being at the centre of the circle, and  $x', y'$ , and  $x'', y''$ , coordinates of the points where the given lines cut the circumference of the given circle. Then we easily obtain:

$$\begin{aligned} xx' + yy' &= r^2 \dots (1); & xx'' + yy'' &= r^2; \dots (2) \\ x'^2 + y'^2 &= r^2 \dots (3); & x''^2 + y''^2 &= r^2; \dots (4) \\ (x' - x'')^2 + (y' - y'')^2 &= (x' - a)^2 + y'^2 + (x'' - a)^2 + y''^2 \\ &\quad - 2 \sqrt{[(x' - a)^2 + y'^2][(x'' - a)^2 + y''^2]} \cos \alpha; \text{ or,} \\ x'x'' + y'y'' - a(x' + x'') + a^2 &= \sqrt{[(r^2 + a^2 - 2ax')(r^2 + a^2 - 2ax'')]} \cos \alpha \dots (5) \end{aligned}$$

Combining (1) and (3),

$$x' = \frac{r}{x^2 + y^2} \left( rx \pm y \sqrt{(x^2 + y^2 - r^2)} \right); \quad y' = \frac{r}{x^2 + y^2} \left( ry \mp x \sqrt{(x^2 + y^2 - r^2)} \right).$$

Combining (2) and (4),

$$x'' = \frac{r}{x^2 + y^2} (rx \mp y \sqrt{x^2 + y^2 - r^2}); \quad y'' = \frac{r}{x^2 + y^2} (ry \pm x \sqrt{x^2 + y^2 - r^2}).$$

Substituting these values in (5), and reducing,

$$[2r^4 - (r^2 - a^2)(x^2 + y^2) - 2ar^2x]^2 = [(r^2 + a^2)^2(x^2 + y^2)^2 - 4ar^2(r^2 + a^2)(x^2 + y^2)x + 4a^2r^2(r^2 - y^2)(x^2 + y^2)] \cos^2 \alpha;$$

or, in a better form for discussion,

$$[2r^4 - (r^2 - a^2)(x^2 + y^2) - 2ar^2x]^2 \tan^2 \alpha = 4r^2(x^2 + y^2 - r^2)(ax - r^2)^2.$$

When  $\alpha = 0$ , we have  $x^2 + y^2 - r^2 = 0$ , and  $(ax - r^2)^2 = 0$ , the original circle, and two coincident straight lines. When  $\alpha = 90^\circ$ ,  $[2r^4 - (r^2 - a^2) \times (x^2 + y^2) - 2ar^2x]^2 = 0$ , two coincident circles.

As  $\alpha$  increases from 0, the circle passes into an oval with the horizontal axis the longer, while the two straight lines separate at first into two infinite branches. The right hand branch recedes to infinity and then reappears on the left, forming now with the other line an oval whose vertical axis is the longer. These ovals then approach till, when  $\alpha = 90^\circ$ , they coincide in a circle.

When  $\alpha = 0$ , the equation becomes

$$(x^2 + y^2 - r^2 \operatorname{cosec}^2 \frac{1}{2} \alpha)(x^2 + y^2 - r^2 \sec^2 \frac{1}{2} \alpha) = 0,$$

two circles, coincident when  $\alpha = 90^\circ$ , as should be expected.

When  $\alpha = r$ , the equation becomes  $(x^2 + y^2 - r^2 \sec^2 \alpha)(x - r)^2 = 0$ , a circle and two coincident straight lines.

The locus can be easily constructed for a number of values of  $\alpha$  at the same time, and the interesting changes seen at a glance.

[Dr. Eggers gets

$2(px + qy - r^2) \sqrt{x^2 + y^2 - r^2} = [2(px + qy - r^2) + (r^2 - p^2 - q^2)(x^2 + y^2)] \tan \alpha$ , for the required equation; in which  $p$  and  $q$  are coordinates of the point  $O$ .

Mr. Seitz gets

$$\cos \theta = \frac{r^2}{a\rho} - \frac{(r^2 - a^2)\rho \tan \alpha}{2ar[r \tan \alpha - \sqrt{(\rho^2 - r^2)}]},$$

for the polar equation of the curve; where  $\rho$  is the radius vector and  $\theta$  the angle between the radius vector and a diameter through  $O$ .]

102. "Prove that in every triangle the square of the sum of the squares of the sides exceeds double the sum of their fourth powers."

SOLUTION BY E. B. SEITZ, GREENVILLE, OHIO.


LET  $a, b, c$  be the sides. Then  $a + b - c > 0$ ,  $a + c - b > 0$ ,  $b + c - a > 0$ , and  $a + b + c > 0$ . By multiplication,  $-a^4 - b^4 - c^4 + 2a^2b^2 + 2a^2c^2 + 2b^2c^2 > 0$ . Adding to both memb's  $2(a^4 + b^4 + c^4)$ , we have  $(a^2 + b^2 + c^2)^2 > 2(a^4 + b^4 + c^4)$ .

103. "Through two given points draw a circle bisecting the circumference of a given circle."

SOLUTION BY HENRY HEATON, DESMOINES, IOWA.

LET  $A$  and  $B$  represent the given points and  $GFE$  the given circle.

Through  $A$  and  $B$  describe any circle  $ABEF$ , intersecting the given circle in  $E$  and  $F$ . Draw  $EF$  and prolong it until it meets  $AB$  produced in  $H$ . Through the center  $C$ , draw  $HG$  cutting the given circle in  $D$  and  $G$ ; then will  $D$  and  $G$  be points in the required circle. For  $AH \times BH = FH \times EH = GH \times DH$ ; and if a circle be passed through the points  $A, B$  and  $D$  it will cut  $HG$ , or  $HG$  produced, at some point  $I$ ; and we have  $AH \times BH = IH \times DH = GH \times DH$ . Hence  $I$  coincides with  $G$ , and the circle  $ABD$  passes through  $G$ .



[The solutions by Dr. Eggers, Prof. Johnson, Prof. M. C. Stevens and Mr. Seitz are substantially the same as the above. Prof. Stevens' solution was received too late to be included in our notice of solutions, published in No. 2. We have, also, an elegant solution of this question by Miss Ladd which should have been included with the notices in No. 2, but was omitted through an oversight.]

104. "Show that  $x^3 + 1 = y^2$  is possible for the values  $x=0, -1$ , and  $1$ , only."

SOLUTION BY PROF. A. B. EVANS, LOCKPORT, NEW YORK.

IF  $x^3 + 1 = y^2$  we may put  $x = m - 1$ ; then

$$x^3 + 1 = m^3 - 3m^2 + 3m = \square. \dots \dots \dots (1)$$

It is evident that (1) must be divisible by  $m^2$ ; put, therefore,

$$m^3 - 3m^2 + 3m = a^2 m^2; \text{ then } m^2 - (a^2 + 3)m = -3, \dots (2)$$

and

$$m = 0. \dots \dots \dots (3)$$

From equation (2),  $m = \frac{1}{2}(a^2 + 3) \pm \frac{1}{2}\sqrt{(a^2 + 3)^2 - 12}$ . Hence  $(a^2 + 3)^2 - 12 = \square$ .

Now  $a^2 + 3 = 4$  is the only value of  $(a^2 + 3)$  in integers that will satisfy this condition;  $\therefore a = \pm 1, m = 2 \pm 1 = 3$  or  $1, x = m - 1 = 2$  or  $0$ . From equation (3)  $x = -1$ .

[Dr. Nelson requests us to state that problems 104 and 105 were contributed as selected and not as original. The above solution shows that the value  $x = 1$ , &c., as announced, should have been  $x = 0, -1$ , and  $2$ .]

